

Procter & Gamble

PHARMACEUTICALS

Mail: The Procter & Gamble Company
Health Care Research Center
P.O. Box 8006
Mason, Ohio 45040-8006

Shipping: The Procter & Gamble Company
Health Care Research Center
8700 Mason-Montgomery Road
Mason, Ohio 45040-9462

3 June 2002

Dockets Management Branch
HFA-305
Food and Drug Administration
5630 Fishers Lane
Room 1061
Rockville, MD 20852

Re: Docket No. 02D-0095
Draft Guidance for Industry on Exposure-Response Relationships: Study Design, Data Analysis, and Regulatory Applications

Dear Sir or Madam:

Reference is made to the Draft Guidance for Industry on Exposure-Response Relationships: Study Design, Data Analysis, and Regulatory Applications. Submitted herewith, are comments from Procter & Gamble Pharmaceuticals regarding the draft guidance document. We appreciate the opportunity to respond to the Agency's request for comments.

Comments

Section IV. B. Using observed concentration versus concentration-controlled trials. One would agree that if everyone is administered the same dose, the ability to assess the influence of different covariates on the exposure /response relationships may be difficult due to confounding effects (e.g. renal impairment and higher concentrations). However, the discussion should also mention that if dosage adjustments are implemented into Phase 2/3 for appropriate subgroups, the ability to assess the influence of these subgroups (e.g. renal impairment) may be significantly improved.

Line 413 – Please clarify your use of "rigor" which is always required.

Lines 422/423. Randomization is critical at all stages, not just for those studies intended to support regulatory decisions.

Table I. Points to Consider in Study Design and Exposure-Response Study Analysis. Consider the addition of blinding as a point to consider for all studies, especially if it includes subjective assessment or assessors.

Section V.C.1.e. In addition to changes in plasma protein binding between various diseases states, time-dependent binding should also be mentioned since it may also complicate the analysis of exposure/response analysis.

Lines 514/521. AUC should be replaced by Cavg. It maintains consistency between the various types of exposure (all concentrations) used to assess relationships. More importantly, it allows one to appropriately analyze data from a study where the dosing interval is varied, in an attempt to reduce the

02D-0095

C5

correlation between the various measures of exposure. It is unclear how one would use AUC_{τ} in the analysis when the dosing interval is varied without standardizing to the dosing interval (i.e., C_{avg}).

Thank you again for the opportunity to provide comments. If you have any questions regarding the above, please contact the undersigned by telephone at 513.622.5278, or by facsimile at 513.622.5363.

Sincerely,

A handwritten signature in black ink that reads "Wendy Sauber". The signature is written in a cursive, flowing style.

Wendy M. Sauber
Section Head, U.S. Regulatory Affairs

MICHAEL D SHILLEN
PROCTOR & GAMBLE MCRC
8700 MASON MONTGOMERY RD.
MASON OH 45040

(513)622-1376

TO: ROOM 1061
FOOD & DRUG ADMIN.
5630 FISHERS LANE
HFA-305
ROCKVILLE

SHIP DATE: 03JUN02
ACC # 175926763

ACTUAL WGT: 1 LBS SCALE

4987 0859 5243

FedEx

MD 20857

REF: 6758HC2961

PRIORITY OVERNIGHT

TUE::

cad # 0626125 03JUN02

TRK# 4987 0859 5243

FORM
0201

Deliver by:
04JUN02

20857 -MD-US

IAD AA

NZ GAIA



The World On Time